arXiv:1704.00757v1 [math.CV] 3 Apr 2017

NORMING SETS ON A COMPACT COMPLEX MANIFOLD

TANAUSÚ AGUILAR-HERNÁNDEZ

Abstract. We describe the norming sets for the space of global holomorphic sections to a k-power of a positive holomorphic line bundle on a compact complex manifold X. We characterize in metric terms the sequence of measurable subsets $\{G_k\}_k$ of X such that there is a constant $C > 0$ where

$$\|s\|^2 \leq C \int_{G_k} |s(z)|^2 \, dV(z)$$

for every $s \in H^0(L^k)$ and for all $k \in \mathbb{N}$.

1. Introduction

Let $\mathcal{L}$ be a certain space of functions. The problem lies in giving a metric characterization of the set G that verifies that the integral over G is comparable to the norm for every function of $\mathcal{L}$. As we will see below this problem has already proved for some space of functions.

In [LS74] it appears the proof of the Logvinenko-Sereda theorem for functions of the Paley-Wiener space PW_K for a fixed K:

Theorem 1.1. *For a measurable subset $G \subset \mathbb{R}^n$ the following are equivalent:*

(1) *There is a constant $C > 0$ such that*

$$\int_{\mathbb{R}^n} |f|^2 \, dm \leq C \int_G |f|^2 \, dm$$

for every $f \in PW_K$. We will say that G is a norming set for the Paley-Wiener space.

(2) *There is a cube $Q \subset \mathbb{R}^n$ ans a constant $\gamma > 0$ such that*

$$m((Q + x) \cap G) \geq \gamma$$

for all $x \in \mathbb{R}^n$. We will say that G is relatively dense in $\mathbb{R}^n$.

See also [HJ94, pag 112-115].

The previous theorem has been extended to functions in the Bergman space A^p in the unit disk $\mathbb{D}$ in [Lue81]:

Theorem 1.2. *For a measurable subset $G \subset \mathbb{D}$ the following are equivalent:*

(1) *There is a constant $C > 0$ such that*

$$\int_{\mathbb{D}} |f|^p \, dm \leq C \int_G |f|^p \, dm$$

for every $f \in A^p$. We will say that G is a norming set for the Bergman space.

(2) *There is a constant* $\delta > 0$ *and radius* $R \in (0,1)$ *such that*

$$m(G \cap D(a,R)) > \delta m(D(a,R))$$

for all $a \in \mathbb{D}$. *We will say that* G *is* relatively dense in the disc $\mathbb{D}$.

Remark 1.3. In [Lue81] a different geometry than usual will be used. The disks $D(a,R)$ will be of the form

$$D(a,R) = \{z \in \mathbb{D} \ : \ |z-a| < R(1-|a|)\},$$

where $a \in \mathbb{D}$ and $R \in (0,1)$.

It should be noted that using similar arguments to those used in [Lue81] one can prove the Logvinenko-Sereda theorem for the classical Fock space $\mathcal{F}^2_{2|z|^2}(\mathbb{C}^n)$:

Theorem 1.4. *For a a measurable subset* $G \subset \mathbb{C}^n$ *the following are equivalent:*

(1) *There is a constant* $C > 0$ *such that*

$$\int_{\mathbb{C}^n} |f|^2 e^{-2|z|^2} \, dm(z) \leq C \int_G |f|^2 e^{-2|z|^2} \, dm(z)$$

for every $f \in \mathcal{F}^2_{2|z|^2}(\mathbb{C}^n)$. *We will say that* G *is a* norming set for the classical Fock space.

(2) *There is a constant* $\delta > 0$ *and a radius* $R > 0$ *such that*

$$m(G \cap G(z,R)) \geq \delta m(B(a,R))$$

for all $z \in \mathbb{C}^n$. *We will say that* G *is* relatively dense in $\mathbb{C}^n$.

Our main goal will be to prove the Logvinenko-Sereda theorem for the space of global holomorphic sections to a k-power of a positive line bundle on a compact complex manifold X. Before giving the statement of the theorem, we will present the convenient definitions.

First of all, we consider a compact complex manifold X which will be endowed with a smooth Hermitian metric ω. As we know, this metric induces a distance function $d(x,y)$ on X, which will be used to define the balls, and a volume form V, which will be used to integrate over the manifold.

We assume that the holomorphic line bundle L on a compact complex manifold X is endowed with a smooth hermitian metric ϕ.

We will denote by $H^0(L)$ the space of global holomorphic sections to L. Moreover, we will consider that the line bundle (L,ϕ) is positive.

As ϕ is a Hermitian metric on L, then $\partial\overline{\partial}\phi$ is a globally defined (1,1)-form on X, which is called the curvature form of the metric ϕ. Moreover, the line bundle L with the metric ϕ is called positive if $i\partial\overline{\partial}\phi$ is a positive form. The statement of the Logvinenko-Sereda theorem is the following:

Theorem 1.5. *For a sequence of measurable subsets* $\{G_k\}$ *in* X *the following are equivalent:*

(1) *There is a constant* $C > 0$ *such that for all* $k \in \mathbb{N}$,

$$\int_X |s(x)|^2 \, dV(x) \leq C \int_{G_k} |s(x)|^2 \, dV(x)$$

for every $s \in H^0(L^k)$. *We will say that* G_k *is a* norming set in $H^0(L^k)$.

(2) *There is a constant* $\delta > 0$ *and a radius* R *such that*

$$V\left(G_k \cap B\left(a, \frac{R}{\sqrt{k}}\right)\right) > \delta V\left(B\left(a, \frac{R}{\sqrt{k}}\right)\right)$$

for all $a \in X$ *and* $k \in \mathbb{N}$. *We will say that* G_k *is* relatively dense in X.

Remark 1.6 (Bergman kernel)**.** The space $H^0(L)$ admits a Hilbert space structure endowed with the scalar product

$$\langle u, v\rangle = \int_X \langle u(x), v(x)\rangle, \quad u, v \in H^0(L)$$

where the integration is taken with respect to the volume form V.

The Bergman kernel $\Pi(x, y)$ associated to this space is a section to the line bundle $L \boxtimes \overline{L}$ over the manifold $X \times X$, defined by

$$\Pi(x, y) = \sum_{j=1}^{N} s_j(x) \otimes \overline{s_j(y)}$$

where $s_1, \ldots, s_N$ is an orthonormal basis for $H^0(L)$. Moreover, this definition does not depend on the choice of te orthonormal basis. Notice that $L \boxtimes \overline{L} = \pi_1^*(L) \otimes \pi_2^*(\overline{L})$ where $\pi_i = X \times X \to X$ is the projection onto the i-factor.

The Bergman kernel $\Pi(x, y)$ is in a sense the reproducing kernel for the space $H^0(L)$, satisfying the reproducing formula

$$s(x) = \int_X \langle s(y), \Pi(x, y)\rangle \, dV(y)$$

for $s(y) \in H^0(L)$.

The pointwise norm of the Bergman kernel in symmetric, $|\Pi(x, y)| = |\Pi(y, x)|$. Moreover, it satisfies

$$|\Pi(x, x)| = \int_X |\Pi(x, y)|^2 \, dV(y).$$

Lemma 1.7. *Let* (L, ϕ) *be a positive line bundle. We have the off-diagonal estimate*

$$|\Pi_k(x, y)| \lesssim k^n e^{-c\sqrt{k}\ d(x,y)}$$

where c *is an appropriate positive constant and the diagonal estimate*

$$|\Pi_k(x, x)| \asymp k^n.$$

Moreover, from this we obtain the estimate of the dimension of $H^0(L^k)$

$$\dim H^0(L^k) \asymp k^n.$$

See [Ber03] where this lemma is proved using a method that appears in [Lin01].

2. Previous result

The main goal of the lemma is to prove for each ball the existence of a normalized peak-section, that is, a section such that most of the mass of the function is in such ball. For this we will use the reproducing kernel of $H^0(L^k)$.

Lemma 2.1. *Given $\varepsilon > 0$, there is a radius $R > 0$ such that for all ball $B\left(\xi, \frac{R}{\sqrt{k}}\right)$, there is a section $s = s_{\xi,k} \in H^0(L^k)$ such that*

- $\|s\|^2 = \int_X |s(x)|^2 \, dV(x) = 1$,
- $\int_{X\setminus B\left(\xi,\frac{R}{\sqrt{k}}\right)} |s(x)|^2 \, dV(x) < \varepsilon$,

Proof. Given $\varepsilon > 0$. We consider the reproducing kernel of the Hilbert space $H^0(L^k)$

$$\Pi_k(x,y) = \sum_{j=1}^{N} s_j(x) \otimes \overline{s_j(y)}$$

where $s_1, \ldots, s_N$ is an orthonormal basis for $H^0(L^k)$.

If we fix $y \in X$, there is a section $\Phi_{k,y} \in H^0(L^k)$ such that

$$|\Phi_{k,y}(x)| = |\Pi_k(x,y)|, \quad x \in X,$$

by the lemma in [LOC12, pag. 431].

We will consider the section

$$g_{k,y}(x) = \frac{\Phi_{k,y}(x)}{\sqrt{|\Pi_k(y,y)|}}, \quad x \in X,$$

where Π_k denotes the Bergman kernel for the k'th power L^k of the line bundle L.

Let us see the first property.

$$\int_X |g_{k,y}(x)|^2 \, dV(x) = \frac{1}{|\Pi_k(y,y)|} \int_X |\Pi_k(x,y)|^2 \, dV(x) = \frac{|\Pi_k(y,y)|}{|\Pi_k(y,y)|} = 1.$$

Now, we check the second property

$$\begin{aligned}
&\int_{X\setminus B\left(y,\frac{R}{\sqrt{k}}\right)} |g_{k,y}(x)|^2 \, dV(x) = \frac{1}{|\Pi_k(y,y)|} \int_{X\setminus B\left(y,\frac{R}{\sqrt{k}}\right)} |\Pi_k(x,y)|^2 \, dV(x) \\
&= \frac{1}{|\Pi_k(y,y)|} \int_0^{\infty} V\left(\{x : |\Pi_k(x,y)| > \lambda\} \setminus B\left(y, \frac{R}{\sqrt{k}}\right)\right) 2\lambda \, d\lambda \\
&\leq \frac{1}{|\Pi_k(y,y)|} \int_0^{k^n D} V\left(\left\{x : De^{-c\sqrt{k}\ d(x,y)} \geq k^{-n}\lambda\right\} \setminus B\left(y, \frac{R}{\sqrt{k}}\right)\right) 2\lambda \, d\lambda.
\end{aligned}$$

where we use that $|\Pi_k(x,y)| \leq k^n D e^{-c\sqrt{k}\ d(x,y)}$ for some constants $c, D > 0$, as we see in Lemma 1.7. Applying the change of variable $\lambda = k^n D e^{-cu}$.

$$\begin{aligned}
&\frac{1}{|\Pi_k(y,y)|}\int_0^\infty V\left(B\left(y,\frac{u}{\sqrt{k}}\right)\setminus B\left(y,\frac{R}{\sqrt{k}}\right)\right)(2k^nDe^{-cu})|-ck^nDe^{-cu}|\ du\\
&\lesssim \frac{1}{k^n}\int_R^\infty \left(\frac{u}{\sqrt{k}}\right)^{2n}(2k^nDe^{-cu})|-ck^nDe^{-cu}|\ du\\
&\lesssim \int_R^\infty u^{2n}e^{-2cu}\ du = e^{-2cR}\int_0^\infty(\mu+R)^{2n}e^{-2c\mu}\ d\mu\\
&= e^{-2cR}\sum_{j=0}^{2n}\binom{2n}{k}R^{2n-j}\int_0^\infty \mu^j e^{-2c\mu}\ d\mu\\
&\lesssim e^{-2cR}\sum_{j=0}^{2n}\binom{2n}{k}\frac{R^{2n-j}}{(2c)^j}j! \lesssim e^{-2cR}\sum_{j=0}^{2n}\binom{2n}{k}(2cR)^{2n-j} = e^{-2cR}(1+2cR)^{2n}.
\end{aligned}$$

Finally, for R large enough we obtain that

$$\int_{X\setminus B\left(y,\frac{R}{\sqrt{k}}\right)}|g_{k,y}(x)|^2\ dV(x) < \varepsilon.$$

□

Remark 2.2. Notice that taking the same section of the previous lemma and using the Cauchy-Schwarz inequality we obtain that

$$|s(x)|^2 = |\langle s(y), \Pi_k(x,y)\rangle|^2 \leq \|s\|^2|\Pi(x,x)| \lesssim k^n.$$

3. Main results

Theorem 3.1. *For a sequence of measurable subset $\{G_k\}_k$ in X the following are equivalent:*

(1) *There is a constant $C > 0$ such that for all $k \in \mathbb{N}$,*

$$\int_X |s(x)|^2\ dV(x) \leq C\int_{G_k}|s(x)|^2\ dV(x)$$

for every $s \in H^0(L^k)$. We will say that G_k is a norming set of $H^0(L^k)$.

(2) *There is a constant $\delta > 0$ and a radius R such that*

$$V\left(G_k \cap B\left(a,\frac{R}{\sqrt{k}}\right)\right) > \delta V\left(B\left(a,\frac{R}{\sqrt{k}}\right)\right)$$

for all $a \in X$ and $k \in \mathbb{N}$. We will say that G_k is relatively dense in X.

Proof. The proof that (1) implies (2) is the easiest. In this proof, we consider a section with the properties of Lemma 2.1.

So, given $\varepsilon \leq 1/2C$ and applying the Lemma 2.1 there is a radius R such that for all balls $B\left(\xi,\frac{R}{\sqrt{k}}\right)$ there is a section s verifying the properties of the lemma. Hence,

applying the Remark 2.2 we obtain

$$\frac{V\left(G_k \cap B\left(\xi, \frac{R}{\sqrt{k}}\right)\right)}{V\left(B\left(\xi, \frac{R}{\sqrt{k}}\right)\right)} \gtrsim \frac{1}{R^{2n}} \int_{G_k \cap B\left(\xi, \frac{R}{\sqrt{k}}\right)} |s^*(x)|^2 \, dV(x)$$

and using the other properties with (2) we have

$$\begin{aligned}\frac{V\left(G_k \cap B\left(\xi, \frac{R}{\sqrt{k}}\right)\right)}{V\left(B\left(\xi, \frac{R}{\sqrt{k}}\right)\right)} &\gtrsim \frac{1}{R^{2n}} \left(\int_G |s^*(x)|^2 \, dV(x) - \int_{X \setminus B\left(\xi, \frac{R}{\sqrt{k}}\right)} |s^*(x)|^2 \, dV(x) \right) \\ &\geq \frac{1}{R^{2n}} \left(\frac{1}{C} - \varepsilon \right) \geq \frac{1}{2CR^{2n}}.\end{aligned}$$

Therefore, we have proved that (1) implies (2).

The proof that (2) implies (1) is the difficult one. We will use a similar argument as in [MOC08, Chapter 4]. We will partition X in two pieces. The first piece, denoted by $\mathcal{A}$, are the points where the sections is much smaller that its average and the second is the complementary $X \setminus \mathcal{A}$. The following lemma proves that the integral over $\mathcal{A}$ is irrelevant as most of the mass is carried by $X \setminus \mathcal{A}$.

Lemma 3.2. *Let $\varepsilon > 0$ and $s \in H^0(L^k)$. Define the set*

$$\mathcal{A} = \left\{ a \in X \mid |s(a)|^2 < \frac{\varepsilon}{V(B(a, R/\sqrt{k}))} \int_{B(a,R/\sqrt{k})} |s(z)|^2 \, dV(z) \right\}.$$

Then there is a constant C depending on R and k such that

$$\int_{\mathcal{A}} |s(z)|^2 \, dV(z) < C\varepsilon \int_X |s(z)|^2 \, dV(z).$$

Proof. For $\xi \in \mathcal{A}$ we have

$$|s(a)|^2 < \frac{\varepsilon}{V(B(a, R/\sqrt{k}))} \int_{B(a,R/\sqrt{k})} |s(z)|^2 \, dV(z) = \varepsilon \int_X |s(z)|^2 \frac{\chi_{B(a,R/\sqrt{k})}(z)}{V(B(a, R/\sqrt{k}))} \, dV(z).$$

Integrating respect to a and applying the Fubini's theorem

$$\begin{aligned}\int_{\mathcal{A}} |s(a)|^2 \, dV(a) &< \varepsilon \int_X |s(z)|^2 \left(\int_{\mathcal{A}} \frac{\chi_{B(a,R/\sqrt{k})}(z)}{V(B(a, R/\sqrt{k}))} \, dV(a) \right) \, dV(z) \\ &\leq \frac{V(B(a, R/\sqrt{k}))}{V(B(a, R/\sqrt{k}))} \varepsilon \int_X |s(z)|^2 \, dV(z) = \varepsilon \int_X |s(z)|^2 \, dV(z).\end{aligned}$$

Therefore, we obtain

$$\int_{\mathcal{A}} |s(a)|^2 \, dV(a) \leq \varepsilon \int_X |s(z)|^2 \, dV(z).$$

□

Let $\mathcal{F} = X \setminus \mathcal{A} = \left\{a \in X \ : \ |s(a)|^2 \geq \frac{\varepsilon}{V(B(a,R/\sqrt{k}))} \int_{B(a,R/\sqrt{k})} |s(z)|^2 \ dV(z)\right\}$. If we choose ε such that $\varepsilon C < 1/2$ we have

$$\int_X |s(z)|^2 \ dV(z) < 2 \int_{\mathcal{F}} |s(z)|^2 \ dV(z)$$

since

$$\begin{aligned}\int_X |s(z)|^2 \ dV(z) &= \int_{\mathcal{F}} |s(z)|^2 \ dV(z) + \int_{\mathcal{A}} |s(z)|^2 \ dV(z) \\ &< \int_{\mathcal{F}} |s(z)|^2 \ dV(z) + C\varepsilon \int_X |s(z)|^2 \ dV(z) \\ &< \int_{\mathcal{F}} |s(z)|^2 \ dV(z) + \frac{1}{2} \int_X |s(z)|^2 \ dV(z).\end{aligned}$$

Thus this is enough to show that

$$\int_{\mathcal{F}} |s(z)|^2 \ dV(z) \lesssim \int_{G_k} |s(z)|^2 \ dV(z).$$

All we need to prove is the existence of a constant $C > 0$ such that for all $w \in \mathcal{F}$

$$|s(w)|^2 \leq \frac{C}{V\left(B\left(w, \frac{R}{\sqrt{k}}\right)\right)} \int_{B\left(w,\frac{R}{\sqrt{k}}\right)\cap G_k} |s(z)|^2 \ dV(z). \tag{3.1}$$

Indeed, if this is the case then

$$\begin{aligned}\int_{\mathcal{F}} |s(w)|^2 \ dV(w) &\leq C \int_{G_k} |s(z)|^2 \left(\int_{\mathcal{F}} \frac{\chi_{B\left(w,\frac{R}{\sqrt{k}}\right)}(z)}{V\left(B\left(w, \frac{R}{\sqrt{k}}\right)\right)} \ dV(w) \right) \ dV(z) \\ &\leq C \int_{G_k} |s(z)|^2 \ dV(z).\end{aligned}$$

To prove (3.1) we argue by contradiction. If (3.1) is false there are for any $q \in \mathbb{N}$ sections $s_q \in H^0(L^{k_q})$ and $w_q \in \mathcal{F}$ such that

$$|s_q(w_q)|^2 > \frac{q}{V\left(B\left(w_q, \frac{R}{\sqrt{k_q}}\right)\right)} \int_{B(w_q,\frac{R}{\sqrt{k_q}})\cap G_{k_q}} |s_q(z)|^2 \ dV(z) \tag{3.2}$$

By compactness of X we can choose a convergent subsequence $\{w_q^*\}$ of w_q to some $w^* \in X$ such that there is a local chart (U, φ) where

$$B\left(w_q^*, \frac{R}{\sqrt{k_q}}\right) \subset U, \ \forall q \in \mathbb{N}$$

and $\varphi(w^*) = 0$.

By the positivity of $i\partial\overline{\partial}\phi$, we can make a linear change of variable so that $i\partial\overline{\partial}\phi(\varphi^{-1}(0)) = i\partial\overline{\partial}|z|^2$. So if we consider the Taylor series of ϕ in the local ball $B(0,\tau)$ we obtain

$$\phi(\varphi^{-1}(z)) = h(z) + |z|^2 + q(z)$$

where h is a pluriharmonic polynomial of degree less than or equal to 2, $h(0) = \phi(\varphi^{-1}(0))$ and $q(z) = o(|z|^2)$. Hence, we have that

$$|2\phi(\varphi^{-1}(z)) - 2|z|^2 - 2h(z)| \leq c(\tau)\tau^2$$

in $B(0,\tau)$ for nondecreasing continuous function $c(\tau)$ such that $c(0) = 0$. Notice that we can use the local chart $(\varphi^{-1}(B(0,\tau*)),\varphi)$ instead since $\exists \nu \in \mathbb{N}$, $\forall q \geq \nu$: $B\left(w_q^*, \frac{R}{\sqrt{k_q}}\right) \subset \varphi^{-1}(B(0,\tau^*))$ and $\tau^* \leq \tau$.

Now we consider the sequence of holomorphic functions

$$G_q(z) = f_q(\varphi^{-1}(z))e^{-\frac{k}{2}H(z)}, \quad q \in \mathbb{N}$$

where $\Re(H) = h$ and $|s_q|^2 = |f_q|^2 e^{-k\phi}$.

Next we need obtain that

$$|s_q(w_q^*)|^2 \lesssim |G_q(\eta_q)|^2$$

where $\varphi(w_q^*) = \eta_q$,

$$|s_q(\varphi^{-1}(z))|^2 \gtrsim G_q(z)|^2$$

for a certain ball and

$$|G_q(\lambda_q)|^2 \gtrsim \frac{q}{m(B(\eta_q,\delta_q))}\int_{B(\eta_q,\delta_q)\cap\varphi(E_{k_q})} |G_q(z)|^2 \, dm(z). \tag{3.3}$$

First, for the sequence $\{w_q^*\}$ we have

$$|s_q(w_q^*)|^2 = |f_q(\varphi^{-1}(\eta_q))|^2 e^{-k\phi(\varphi^{-1}(\eta_q))} \leq |f_q(\varphi^{-1}(\eta_q))|^2 e^{-kh(\eta_q)} e^{\frac{k}{2}c(\tau^*)\tau^*} \lesssim |G_q(\eta_q)|^2.$$

Furthermore for $z \in B(0,\tau^*)$ we obtain

$$|G_q(z)|^2 = |f_q(\varphi^{-1}(z))|^2 e^{-kh(z)} \lesssim |s_q(\varphi^{-1}(z))|^2, \quad z \in B(0,\tau^*).$$

Finally we obtain the inequality that we need

$$\begin{aligned}
|G_q(\eta_q)|^2 \gtrsim |s_q(w_q^*)|^2 &\gtrsim \frac{q}{V\left(B\left(w_q^*, \frac{R}{\sqrt{k_q}}\right)\right)} \int_{B\left(w_q^*, \frac{R}{\sqrt{k_q}}\right)\cap E_{k_q}} |s_q(z)|^2 \, dV(z) \\
&\gtrsim \frac{q}{V\left(B\left(w_q^*, \frac{R}{\sqrt{k_q}}\right)\right)} \int_{\varphi\left(B\left(w_q^*, \frac{R}{\sqrt{k_q}}\right)\cap E_{k_q}\right)} |f_q(\varphi^{-1}(z))|^2 \, e^{-k\phi(\varphi^{-1}(z))} \, dm(z) \\
&\gtrsim \frac{q}{m(B(\eta_q,\delta_{k_q}))} \int_{B\left(\eta_q,\delta_{k_q}\right)\cap E_{k_q}^*} |G_q(z)|^2 \, dm(z)
\end{aligned}$$

where $E_{k_q}^* = \varphi(E_{k_q})$ and there is a ball $B(\eta_{k_q},\delta_{k_q}) \subset \varphi(w_q^*, R/\sqrt{k_q})$ such that $m(B(\eta_q,\delta_{k_q})) \asymp V\left(B\left(w_q^*, \frac{R}{\sqrt{k_q}}\right)\right)$, because as $\varphi, \varphi^{-1} \in \mathcal{C}^1$ we have that φ is bi-Lipschitz.

By means of a dilatation and a translation we send η_q to the origin of $\mathbb{C}^n$, the ball $B(\eta_q,\delta_{k_n})$ to $B(0,1) \subset \mathbb{C}^n$ and the set $E_{k_q}^*$ to E_q^*. We will denote the set $\overline{B(0,1)}\cap E_q^*$ by H_q.

Moreover, multiplying by a constant we can assume that

$$\int_{B(0,1)} |\tilde{G}_q(z)|^2 \, dm = 1.$$

The subharmoniticity of $|\tilde{G}_q|^2$ and the fact that $w_q^* \in \mathcal{F}$ tells us that

$$\varepsilon \lesssim |\tilde{G}_q(0)|^2 \lesssim 1.$$

Notice that we can modify the inequality of the definition of $\mathcal{F}$ in the same way as the expression (3.3).

And this property together with (3.3) yields

$$\frac{1}{q} \gtrsim \int_{H_q} |\tilde{G}_q(u)|^2 \, dm(u). \tag{3.4}$$

We have that $\{\tilde{G}_q\}$ is a locally bounded sequence of holomorphic functions defined in $B(0,1)$ and therefore using Montel theorem there exist a subsequence converging locally uniformly on $B(0,1)$ to some holomorphic function G.

We observe that the relative dense hypothesis yields

$$\inf_n m(H_n) > 0.$$

The Helly selection theorem guaranties the existence of a weak-$*$limit τ of a subsequence of $\tau_q = \chi_{H_q} m$ such that $\tau \not\equiv 0$. Condition (3.4) implies that $\tilde{G} = 0$ τ-a.e. and therefore $supp\tau \subset \{\tilde{G} = 0\}$.

We want to show that $supp\tau$ cannot lie on a complex $(n-1)$-dimensional submanifold $S \subset \mathbb{C}^n$. We argue by contradiction.

By definition of τ_n we have $\tau_n(B(x,r)) \leq m(B(x,r)) \lesssim r^{2n}$. As this inequality is true for all τ_q, the same holds for τ. But, using the Frostman lemma we obtain that $\dim_{Haus}(supp\tau) \geq 2n > 0$.

Therefore, we have proved that (2) implies (1).

□

Now, we will generalize the previous sampling problem using a sequence of measures $\{\mu_k\}_k$ of $\mathfrak{M}(X)$ instead of the particular sequence of measures $\{\chi_{G_k} V\}_k$ of $\mathfrak{M}(X)$, where $\{G_k\}_k$ are measurable sets. We will solve part of this problem, the remaining part continues being an open problem.

Theorem 3.3. *For a sequence of measures $\{\mu_k\}$ the following are equivalent:*

(1) *There exists a constant $C_1 > 0$ such that for all $k \in \mathbb{N}$,*

$$\int_X |s(z)|^2 \, d\mu_k(z) \leq C_1 \int_X |s(z)|^2 \, dV(z)$$

for every $s \in H^0(L^k)$.

(2) *There exists a constant $C_2 > 0$ such that for all $k \in \mathbb{N}$,*

$$T[\mu_k](z) = \int_X \left| \frac{\Pi_k(w,z)}{\sqrt{|\Pi_k(z,z)|}} \right|^2 d\mu_k(w) \leq C_2$$

for every $z \in X$, that is, the sequence of Berezin transforms $\{T[\mu_k]\}_k$ are uniformly bounded in X.

(3) *There is a constant* $C_3 > 0$ *such that*

$$\mu_k(B(z, 1/\sqrt{k})) \leq \frac{C_2}{k^n}, \qquad z \in X.$$

Proof. The proof that (1) implies (2) is trivial because $\Pi_k(z, w) \in H^0(L^k)$ for each $w \in X$ and $k \in \mathbb{N}$.

To prove that (2) implies (3) is enough the following lemma.

Lemma 3.4. *There are constants* $\varepsilon > 0$, $M > 0$ *and* $k_0 \in \mathbb{N}$ *such that*

$$\left| \frac{\Pi_k(w, z)}{\sqrt{|\Pi_k(z, z)|}} \right|^2 \geq Mk^n > 0 \tag{3.5}$$

for every $w \in B(z, \varepsilon/\sqrt{k})$ *and* $k > k_0$.

Proof. To prove (3.5) we use that

$$\left| \frac{\Pi_k(z, z)}{\sqrt{|\Pi_k(z, z)|}} \right|^2 - \left| \frac{\Pi_k(w, z)}{\sqrt{|\Pi_k(z, z)|}} \right|^2 = \int_0^1 \frac{d}{dt} \left(\left| \frac{\Pi_k(w - (z - w)t, z)}{\sqrt{|\Pi_k(z, z)|}} \right|^2 \right) dt$$

and by the bound of the derivatives in [LOC12, pag. 434]

$$\left| \frac{\partial}{\partial \xi_j} \left| \frac{\Pi_k(w, z)}{\sqrt{|\Pi_k(z, z)|}} \right|^2 \right| \leq C\sqrt{k} \sup_{\gamma \in U_k(w)} \left| \frac{\Pi_k(\gamma, z)}{\sqrt{|\Pi_k(z, z)|}} \right|^2$$

where

$$U_k(w) = \{x \in X \ \ |z_j - w_j| < 1/\sqrt{k}, (j = 1, \ldots, n)\},$$

together with the sub-mean value property in [LOC12, pag. 432], we obtain that

$$\left| \frac{\Pi_k(z, z)}{\sqrt{|\Pi_k(z, z)|}} \right|^2 - \left| \frac{\Pi_k(w, z)}{\sqrt{|\Pi_k(z, z)|}} \right|^2 \leq M' k^{\frac{2n+1}{2}} \sqrt{n} |z - w|.$$

By the diagonal estimate of Lemma 1.7 we have

$$\left| \frac{\Pi_k(w, z)}{\sqrt{|\Pi_k(z, z)|}} \right|^2 \geq Dk^n - M' k^{\frac{2n+1}{2}} \sqrt{n} |z - w|,$$

where $D > 0$ is a certain constant.

Then, if we consider the ball $B(z, \varepsilon/\sqrt{k})$ where $\varepsilon < \frac{D}{2M'\sqrt{n}}$ we have

$$\left| \frac{\Pi_k(w, z)}{\sqrt{|\Pi_k(z, z)|}} \right|^2 \geq (D - M'\varepsilon\sqrt{n})k^n > \frac{Dk^n}{2} > 0$$

for $w \in B(z, \varepsilon/\sqrt{k})$. □

Applying Lemma 3.4 we have that there are constants $\varepsilon > 0$, $M > 0$ and $k_0 \in \mathbb{N}$ such that

$$Mk^n \mu_k(B(z, \varepsilon/\sqrt{k})) \leq \int_{B(z,\varepsilon/\sqrt{k})} \left| \frac{\Pi_k(w,z)}{\sqrt{|\Pi_k(z,z)|}} \right|^2 d\mu_k(w) \leq T[\mu_k](z) \leq C_2$$

for every $z \in X$ and $k > k_0$.

We take a covering of the ball $B(z, 1/\sqrt{k})$ by a collection of balls $B(z_j, \varepsilon/5\sqrt{k})$, $z_j \in B(z, 1/\sqrt{k})$ and applying the Vitali covering lemma we obtain a sub-collection of these balls which are disjoint such that

$$\cup B(z_j, \varepsilon/5\sqrt{k}) \subset \cup B(z_{j_q}, \varepsilon/\sqrt{k}).$$

Moreover, as

$$B(z_{j_q}, \varepsilon/\sqrt{k}) \subset B(z, 2/\sqrt{k})$$

by means of the volume we have that

$$N \cdot \left(\frac{\varepsilon}{\sqrt{k}}\right)^{2n} \lesssim \left(\frac{2}{\sqrt{k}}\right)^{2n},$$

that is, $N \lesssim \left(\frac{2}{\varepsilon}\right)^{2n}$ where N is the number of balls of the sub-collection. Therefore, we conclude

$$\mu_k(B(z, 1/\sqrt{k})) \lesssim \left(\frac{2}{\varepsilon}\right)^{2n} m_k(B(z_{j_q}), \varepsilon/\sqrt{k}) \lesssim \frac{1}{k^n}.$$

Notice that for $k \leq k_0$ (3) holds immediately since $\mu_k(B(z, 1/\sqrt{k}))$ are bounded by $\mu_k(X)$.

All we need to prove that (3) implies (1) is the existence of a constant $Q > 0$ such that for all $w \in X$

$$|s(w)|^2 \leq Qk^n \int_{B(w,1/\sqrt{k})} |s(z)|^2 \, dV(z), \quad w \in X. \tag{3.6}$$

This is proved by the sub-mean value property in [LOC12, pag. 432]. Indeed, if this is the case then

$$\begin{aligned} \int_X |s(w)|^2 \, d\mu_k &\leq Qk^n \int_X |s(z)|^2 \left(\int_X \chi_{B(w,1/\sqrt{k})}(z) \, d\mu_k(w) \right) \, dV(z) \\ &\lesssim \int_X |s(z)|^2 \, dV(z) \end{aligned}$$

for every $s \in H^0(L^k)$.

□

Example 3.5 (Tautological line bundle)**.** We will apply the Theorem 3.1 to the complex projective space $\mathbb{CP}^n$ with the hyperplane bundle $\mathcal{O}(1)$, endowed with the Fubini-Study metric. Notice that the holomorphic sections to $\mathcal{O}(k)$, the k'th power of $\mathcal{O}(1)$, can be identified with the homogeneous polynomials of degree k in homogeneous coordinates.

Using Theorem 3.1 we obtain that for a given sequence of measurable subsets $\{\Omega_k\}_k$ in $\mathbb{C}^n$, we have that the following statements are equivalent:

(1) For all $k \in \mathbb{N}$, it is verified that

$$\int_{\Omega_k} \frac{|p_k(z)|^2}{(1+|z|^2)^k} \frac{dV(z)}{(1+|z|^2)^{n+1}} \asymp \int_{\mathbb{C}^n} \frac{|p_k(z)|^2}{(1+|z|^2)^k} \frac{dV(z)}{(1+|z|^2)^{n+1}}$$

for every polynomial p_k of degree less or equal than k in n variables.

(2) There is a radius R such that

$$\int_{\Omega_k} \frac{\chi_{H(z,R/\sqrt{k})}(w)}{(1+|w|^2)^{n+1}}\, dV(w) \asymp \left(\frac{R}{\sqrt{k}}\right)^{2n}$$

for all $z \in \mathbb{C}^n$ and $k \in \mathbb{N}$, where

$$H(z, R/\sqrt{k}) = \left\{ w \in \mathbb{C}^n \ : \ |z-w| < \left|\tan\left(\frac{R}{\sqrt{k}}\right)\right| \cdot |1+z\overline{w}| \right\}.$$

Notice that the measure V is the Lebesgue measure on $\mathbb{C}^n$.

Acknowledgement

I would like to express my deep appreciation to Joaquim Ortega Cerdà for his unconditional support and advices to improve in mathematics. In addition, I would like to show my gratitude to my family and friends of Canary Islands for their encouragement and help to fulfil my dreams.

References

[Ber03] B. Berndtsson, *Bergman kernels related to hermitian line bundles over compact complex manifolds*, Explorations in complex and Riemannian geometry, Contemp. Math. **332** (2003), 1–17, MR 2016088 Zbl 1038.32003.

[HJ94] V. Havin and B. Jöricke, *The uncertainty principle in harmonic analysis*, Springer, 1994, ISBN: 978-3-642-78377-7.

[Lin01] N. Lindholm, *Sampling in weighted L^p spaces of entire functions in $\mathbb{C}^n$ and estimates of the bergman kernel*, Journal of Functional Analysis **182** (2001), no. 18, 390–426, doi:10.1006/jfan.2000.3733 MR 1828799 Zbl 1013.32008.

[LOC12] N. Lev and J. Ortega-Cerdà, *Equidistribution estimates for fekete points on complex manifolds*, J. Eur. Math. Soc **18** (2012), no. 2, 425–464, doi:10.4171/JEMS/594.

[LS74] V.N. Logvinenko and J.F. Sereda, *Equivalent norms in spaces of entire functions of exponential type*, Teor. funktsii, funkt. analiz i ich prilozhenia **20** (1974), 62–78.

[Lue81] D.H. Luecking, *Inequalities on Bergman Spaces*, Illinois Journal of Mathematics **25** (1981), no. 1, 1–11, MR 602889 Zbl 0437.30025.

[MOC08] J. Marzo and J. Ortega-Cerdà, *Equivalent norms for polynomials on the sphere*, International Mathematics Research Notices **2008** (2008), doi:10.1093/imrn/rnm154.